

On Raynal's posting in arXiv (nucl-th/0407060)

K. Amos,¹ L. Canton,² G. Pisent,² J. P. Svenne,³ and D. van der Knijff⁴

¹*School of Physics, University of Melbourne, Victoria 3010, Australia*

²*Istituto Nazionale di Fisica Nucleare, sezione di Padova,*

e Dipartimento di Fisica dell'Università di Padova, via Marzolo 8, Padova I-35131, Italia

³*Department of Physics and Astronomy, University of Manitoba,*
and Winnipeg Institute for Theoretical Physics, Winnipeg, Manitoba, Canada R3T 2N2

⁴*Advanced Research Computing, Information Division,*
University of Melbourne, Victoria 3010, Australia

(Dated: February 8, 2008)

With nucl-th/0407060, Jacques Raynal uses the arXiv in a way which does not conform to standard professional practices. His posting contains many statements that are beyond the borders of acceptable scientific disputes, with the scope to defame colleagues by manifestly false or misleading statements. In this comment we reject the three “critiques” expressed by Raynal. 1. The fact that we possibly misquoted our references. 2. The role of the Pauli principle in these kind of calculations. 3. The nature and limits of our coupled-channel potential model. Raynal's postings unfairly detract from the importance of our work, which we published in Nuclear Physics A728, 65 (2003), on a new approach, Multi-Channel-Algebraic-Scattering (MCAS), for coupled-channel calculations. With the MCAS approach we were able to identify systematically all low-energy compound resonances, and to include effectively the Pauli principle in collective, geometrical-type, macroscopic models of multichannel interaction. This represents a clear advantage with respect to the current distribution of the ECIS formulation.

PACS numbers:

Jacques Raynal, in his postings [1, 2], carries criticism of our work and of us personally, and conducts slanderous and unprofessional behaviour, which raises some perplexities on the possible abusive usage of the popular arXiv service. Critiques, even strong criticisms, in science are very healthy and, in general, can be accepted or rejected by confutation. They are part of the scientific method, and they contribute to the development of scientific knowledge. But critiques should adhere to the scientific issue, and provide new information that contributes to clarify the subjects of dispute. Such type of critiques are respectful of the professionalism of the persons and refer only to the scientific debate. A critique should not be used as a means to cast aspersions on colleagues with false and misleading statements, in an attempt to manipulate or misrepresent factual matters for personal interests.

Below we deal with these problems in point fashion. However, the reader should be aware that our disagreements were founded upon a very simple item originally; to wit our use of a reduced form of deformed spin-orbit interaction. Raynal's original objection to our work was that we should have used the fuller derivation in specifying channel interactions in a coupled channels analysis of low-energy neutron ^{12}C scattering; his inference being that our overall approach therefore was totally flawed. In the first posting [1] however, Raynal also initiated a personal attack and presented aspects of our work quite out of context. In our first arXiv response [3] we rejected by confutation his comments, providing our arguments about this very simple item of real debate of a scientific

issue, but we also duly restated the context and scope of our work, which was focused on low-energy compound resonances and specifically on a systematic method of resonance computation and on the inclusion of Pauli principle in geometrical-collective-type approaches.

But the content of Raynal's new posting [2] goes well beyond that, and following discussion with the administrators of *arXiv.org*, we agreed that our present response to these tawdry affairs will be the final one.

1. Raynal asserts that he cannot find in Hodgson's book [4] justification of our claim [3] that Hodgson summarizes the problem of using a deformed optical potential to reproduce a low-energy spectrum of resonances. Raynal is wrong. Pages 426-427 of Hodgson's book section (14.5) are devoted to the problem. It even has the title Resonances with collective excitations. Therein Hodgson describes and presents a very clear physical insight of the problems of using a deformed optical model potential to reproduce the low-energy spectrum of resonances.

2. Raynal looked for discussion of the anti-symmetrization problem in Hodgson's book, and could not find it for scattering. He claims that in [3] we made such use of that reference, and concluded that we do not read the references that we quoted. We never claimed that in Ref. [4] one can find a discussion on the anti-symmetrization problem for scattering. In [3], reference to Hodgson's book was solely related to the summary of the problem of compound resonances and collective excitations; the section 14.5 of Ref. [4].

3. Raynal writes that he cannot find what we claim to be in the other books that we quote in Ref. [3]. Incredi-

bly Raynal also states that, as he could not afford to buy the book by Greiner and Maruhn [5] (nor, apparently, access it in a library), he did not analyze it. How can he then dispute what we claim is contained therein? And in that book particularly the authors note how coupling single particle dynamics with collective (macroscopic-type) degrees of freedom of the target leads to violation of the Pauli exclusion principle. Here we have to be very pedantic: that is precisely the statement to be found on page 297 of Ref. [5], commencing on line 3.

4. Raynal states that he cannot find what we claim to be contained in the book by Mahaux and Weidenmuller [6]. As above, the topic concerns violation of the Pauli principle when coupling single particle dynamics with collective-type degrees of freedom in the target. In that book on page 103, one can find the sentence: “In the papers concerned with ^{13}C as a compound nucleus, the exclusion principle could not be exactly satisfied”. Similar sentences are to be found in pages 113-114.

5. On the basis of the assertions by Raynal of his inability to find that which we attribute to be in the three books, he accuses us of not having read from one or more of the books we quote. Such an accusation calls into question our integrity and we vehemently reject that falsehood.

6. The Mahaux and Weidenmuller book [6] deals with microscopic calculations extensively, and the issue of anti-symmetrization is seriously investigated. Nonetheless in that book, it is also stressed that macroscopic model calculations, using collective (geometrical-deformation) model excitations of the target, violate the Pauli principle. So calculations with one-body type potentials, such as are performed in the ECIS scheme, then have that flaw. The basic calculation we have performed with the method described in Ref. [7] is exactly of that type as well. But in our case the effect of the Pauli principle has been taken into account with the Orthogonalizing Pseudo Potential (OPP) method. That method is applied in our approach by using Eq.(43) in Ref. [7] and it works very well. That success, and the demonstration in Ref. [7] that ignoring the Pauli effect gives much spuriousity, belies Raynal’s insistence that the Pauli principle, though essential in dealing with bound many nucleon systems and (as he rightly observes) is very important at higher energies, somehow has negligible influence on low energy properties.

7. Raynal attempts a stinging criticism of our work by drawing upon a comment made in Ref. [6], namely that the study by Pisent (et al.) is an example of ^{12}C calculations with explicit violation of the exclusion principle. But that is exactly what we resolved in our recent paper [7] on which Pisent is one of the co-authors. The reference in the Mahaux and Weidenmuller book [6] is to work of Pisent made in 1967.

8. Raynal attempts to discredit our claim of the importance of anti-symmetrization by drawing attention to

a paper, on which he is a co-author, as an example of 1 particle- 1 hole $^{16}\text{O}(\gamma, n)$ calculation that “treats anti-symmetrization correctly”. Based on that work Raynal concludes that our claim of an anti-symmetrization problem with the collective-model formulations, such as he has programmed in his code ECIS, is not founded. That is illogical. The fact that Raynal and collaborators made a “microscopic-type” calculation (the definition of which we take as in the Greiner and Maruhn book, p.303), with the required overlap integrals to transform differential equations into integro-differential equations, in no way supports a conclusion that the ECIS formulation, for example, is free from the methodological problems that we have raised. In fact the more appropriate conclusion may well be the opposite with respect to what is claimed by Raynal. To the best of our knowledge, the ECIS formulation is not of a microscopic theory, nor in its current formulation is it suitable for the solution of integro-differential coupled-channel equations when allowance is made for adequate if not complete treatment of the Pauli principle. As well, ECIS is not the basis of the microscopic type calculations made in 1967 by Raynal (et al.). Even for the specific work of Raynal (et al.) so referenced, it is noted in Ref. [6] on p.108, that their (Raynal (et al.)) accord with experimental data at best is only qualitative; something to be expected because “the calculation contains many approximations”. With the above caveat and when the results shown in Fig 6.1 of Ref. [6] are considered, it is clear that these 1967 results of Raynal (et al.) are inconclusive on the role of Pauli effects.

9. Certainly, Raynal is aware of the absolute need to treat anti-symmetrization with nucleon scattering from nuclei, at least with energies of 25 MeV or more. There is a great deal devoted to that in the review of which he is a co-author [8]. In fact therein the results found using Raynal’s other large code, DWBA98, with and without taking anti-symmetrization into account, and with a quite sophisticated microscopic model of the reaction process, are very severe at all energies to pion threshold and beyond. That he should now be dismissive of the role of such at 0 to 5 MeV or more, is not supplemented by any reasonable evidence.

10. In his criticisms [1, 2], Raynal also makes the false claim that our model potential is already symmetric (before symmetrization). Incidentally, had our potential been already symmetric from the start, to symmetrize something already symmetric is not a mistake. It would just be superfluous. But that was not the case in the potential model we considered.

11. Raynal acknowledges that the extra terms in the fuller description of the deformed spin-orbit field plays a role primarily in the asymmetry of the inelastic excitations. With that we concur and noted previously [3] that the effects of the additional terms to the [L S] force have been studied in the past and found to be of minor

import; hardly to be seen to effect with cross sections (inelastic ones) let alone to the elastic cross sections for which it enters in second order. In fact a paper [9] on it shows the effects of this extra term to be quite small in calculating polarizations for all but forward scattering angles; and even there the variations are not particularly large.

12. Raynal contends that with a refitting of parameters of our matrix of potentials, we could have fit data without needing any OPP corrections. It is true that one may find equivalent local potentials to any set of scattering phase shifts. In the review on which Raynal is a coauthor [8], there is a complete chapter on methods and conditions for doing just that. But with phenomenological potential forms the process is "numerical inversion". With such there is no serious constraint of physical nature as to what interaction results. It is also true that by adjusting parameters without constraint one can find interactions that separately fit elastic and inelastic scattering data so one could equally well use Raynal's argument to decry any phenomenological coupled-channel calculation. But one cannot agree with Raynal's argument. A theoretical description should produce sensible results on a variety of observables, and if the excited bound and resonance spectra are so poorly reproduced in their character, one must have first concern with the physics behind the specified model system. Nevertheless, we did make serious attempts at parameter variations in the first instance. We found that even with extreme values for parameters we could never improve the situation with regard to excessive spurious states. The fact is that ignoring the Pauli principle always gives spurious results, while accounting for it leads quickly to sensible results.

13. The three conclusions reached by Raynal are false because: 1) In our posting [3], we have clearly specified what spin-orbit potential we have used and what part of the fuller derivation that is. That fact has never been hidden. 2) We have given exactly what is written and/or can be inferred from content in the three books under scrutiny. We have specified in this letter where such can be found (giving page and line numbers). In light of that, it is simply ridiculous to claim, as Raynal does so sarcastically and unprofessionally, that we did not open one or another of those books. 3) To state that, from Table 1 of Ref. [7], one cannot infer the importance of the Pauli principle is erroneous. In that table, as is made quite clear in the text [7], "With OPP" means that Pauli exclusion has been taken into account, "Without OPP" means that it has not and so is a result of a macroscopic form factor calculation of the same ilk as might be made with ECIS, if such could be used to generate sub-threshold and sharp resonance information. It is clear and evident from the column labeled "Without OPP" in the table of relevance that there are many spurious states in both bound ($E < 0$) and scattering ($E > 0$) regimes. In contrast, by treating the Pauli effects (as we do) to get the

results in the "With OPP" column, the number of sub-threshold bound and scattering states are closely aligned to observed ones and they have the right spin-parities. It is important to note that, without treating the Pauli principle, there are 8 additional spurious excited states and at least two spurious resonances. It is also quite important that accommodating the Pauli principle does not alter results, such as for the higher spin-parity states, for which essentially there is no Pauli blocking since the relevant neutron orbit is unoccupied in the target. That such is the case in our approach, with the OPP corrections, is another validating feature. Instead this fact is quoted erroneously by Raynal as evidence of irrelevance of the Pauli effect.

Finally the concentration by Raynal upon one small element in the total scheme by which we solve the low-energy coupled channels problem is disingenuous. At best it is a smoke screen. Raynal overlooks the utility and adaptability of the method we have developed to analyze low-energy scattering data whatever choice is made for the matrix of interaction potentials. He also overlooks the fact that, despite our choice for the spin-orbit field, that choice was made for convenience and may be changed in future calculations without invalidating the method used with a potentially more convenient form. Indeed with the MCAS approach, a researcher may use whatever he/she opts for that matrix of potentials. Nonetheless the specific choice we have made for the low-energy model gives a matrix of potentials that is rich in structure. The additional phenomenological spin-spin and orbit-orbit terms, taking contributions to second order in the deformation, ensuring that the matrix of potentials is hermitian, lead to results that are overwhelmingly in agreement with mass 13 data. And, more importantly, we have been able to account for the Pauli principle; something that is imperative and not to be trivialized as Raynal has attempted in relation with low-energy scattering.

-
- [1] J. Raynal, nucl-th/0312038 (2003).
 - [2] J. Raynal, nucl-th/0407060 (2004).
 - [3] K. Amos, L. Canton, G. Pisent, J. P. Svenne, and D. van der Knijff, nucl-th/0401055 (2004).
 - [4] P. E. Hodgson, Nuclear Reactions and Nuclear Structure (Clarendon Press, 1971).
 - [5] W. Greiner and J. A. Maruhn, Nuclear models (Springer-Verlag, Berlin, 1996).
 - [6] C. Mahaux and H. A. Weidenmuller, Shell model approach to nuclear reactions (North-Holland, Amsterdam, 1969).
 - [7] K. Amos, L. Canton, G. Pisent, J. P. Svenne, and D. van der Knijff, Nucl. Phys. A728, 65 (2003).
 - [8] K. Amos, P. J. Dortmans, H. V. von Geramb, S. Karataglidis, and J. Raynal, Adv. in Nucl. Phys. 25, 275 (2000).
 - [9] H. Sherif and J.S. Blair, Phys. Lett. **26B**, 489 (1968).